

ALEXANDER POLYNOMIAL OF SEXTICS

MUTSUO OKA

ABSTRACT. Alexander polynomials of sextics are computed in the case of sextics with only simple singularities or sextics of torus type with arbitrary singularities. We will show that for irreducible sextics, there are only 4 possible Alexander polynomials: $(t^2 - t + 1)^j$, $j = 0, 1, 2, 3$. For the computation, we use the method of Esnault-Artal and the classification result in our previous papers.

1. INTRODUCTION

Recall that a sextics is called *of torus type* if it is defined by polynomial $f(x, y) := f_2(x, y)^3 + f_3(x, y)^2$ where $\deg f_i(x, y) = i$ for $i = 2, 3$. Sextics without such a torus expression is called *of non-torus type*. In [10], we have computed the fundamental group of the complement of tame sextics of torus type and we have classified the configurations of the singularities on non-tame sextics of torus type in [9]. In general, the computation of the fundamental groups is more complicated in the case of non-tame torus curves. In this paper, we are interested in another invariant which is called *an Alexander polynomial*. This invariant is weaker than the fundamental groups, but easier to be computed. In fact, there are Zariski pairs with different fundamental group but with the same Alexander polynomial ([7]). The advantage of Alexander polynomial is that it can be computed by the data of the local singularities and the data about their global position in $\mathbf{P}^2$.

It is the purpose of this paper to give a complete atlas of the Alexander polynomials of
(a) sextics of torus type with arbitrary singularities or
(b) sextics with only simple singularities, not necessarily of torus type.

In fact, we will show that there are only four possibilities of Alexander polynomials:

$$1, \quad (t^2 - t + 1), \quad (t^2 - t + 1)^2, \quad (t^2 - t + 1)^3.$$

for irreducible sextics of type (a) or (b). The case $\Delta(t) = (t^2 - t + 1)^3$ corresponds to 9 cuspidal sextics, $\Delta(t) = 1$ corresponds to sextics of non-torus type with $\rho(5) \leq 6$. It is expected that every sextics of non-torus type satisfies the above inequality. The case $\Delta(t) = (t^2 - t + 1)^2$ corresponds to the cases:

$$\Sigma(C) = [8A_2], [8A_2, A_1], [6A_2, E_6], [6A_2, A_5], [6A_2, A_5, A_1], [4A_2, 2A_5], [4A_2, A_5, E_6]$$

and sextics of torus type with configuration $[B_{3,6}, 3A_2]$, $[C_{3,9}, 3A_2]$. For reducible sextics of torus type, we can have further

$$\tilde{\Delta}(t) = (t^2 - t + 1)^2(t^2 + t + 1), (t^2 - t + 1)(t^2 + t + 1),$$

2000 *Mathematics Subject Classification*. 14H30, 14H45, 32S55.

Key words and phrases. Torus curve, Alexander polynomial, fundamental group.

$$(t^2 - t + 1)^2(t^2 + t + 1)(t + 1)^2, (t^2 - t + 1)^4(t^2 + t + 1)^4(t + 1)^4$$

The last case corresponds to a sextic with $B_{6,6}$ and C has 6 line components. See Corollary 12 and Corollary 17 for further detail.

Sextics of non-torus type with non-simple singularities are not considered in this paper.

2. ALEXANDER POLYNOMIAL OF A PLANE CURVE

Let C be an affine curve defined by a polynomial $f(x, y)$ of degree d . We assume that the line at infinity L_∞ and C intersect transversely and we identify $\mathbf{P}^2 - L_\infty \cong \mathbf{C}^2$. Let $\phi : \pi_1(\mathbf{C}^2 - C) \rightarrow \mathbf{Z}$ be the canonical homomorphism induced by the composition of the Hurewicz homomorphism $\psi : \pi_1(\mathbf{C}^2 - C) \rightarrow H_1(\mathbf{C}^2 - C, \mathbf{Z}) \cong \mathbf{Z}^r$ and the summation homomorphism $\mathbf{Z}^r \rightarrow \mathbf{Z}$ where r is the number of irreducible components of C . Let t be a generator of $\mathbf{Z}$.

Let $\tilde{X} \rightarrow \mathbf{C}^2 - C$ be the infinite cyclic covering corresponding to $\text{Ker}\phi$. Then $H_1(\tilde{X}, \mathbf{Q})$ has the structure of Λ module where $\Lambda := \mathbf{Q}[t, t^{-1}]$. Thus we can write as

$$H_1(\tilde{X}, \mathbf{Q}) \cong \mathbf{Q}^{r-1} \oplus \Lambda/\lambda_1(t) \oplus \cdots \oplus \Lambda/\lambda_\nu(t)$$

where $\lambda_i(t)$ is a polynomial with integral coefficients satisfying the properties $\lambda_i(0) \neq 0$ and $\lambda_i | \lambda_{i+1}$ for $i = 1, \dots, \nu - 1$. The *generic Alexander polynomial* $\Delta(t)$ is defined by the product $\Delta(t) = (t - 1)^{r-1} \lambda_1(t) \cdots \lambda_\nu(t)$. Let us call $\tilde{\Delta}(t) := \Delta(t)/(t - 1)^{r-1}$ the *reduced Alexander polynomial*. It does not depend on the choice of the generic line at infinity L_∞ . Let $F(X, Y, Z)$ be the defining homogeneous polynomial of C and let $M = F^{-1}(1) \subset \mathbf{C}^3$ be the Milnor fiber of F . Randell showed in [12] that $\Delta(t)$ can be computed as the characteristic polynomial of the monodromy on the first homology group of the Milnor fibration defined by the homogeneous polynomial $F(X, Y, Z)$. Thus the degree of $\Delta(t)$ is equal to the first Betti number $b_1(M)$. On the other hand, Libgober has proved that the degree of $\tilde{\Delta}(t)$ is equal to the sum $\sum_{j=1}^{d-1} \beta_j$, where β_j is defined by the number of factors $\ell, \ell = 1, \dots, \nu$, such that $\exp(2\pi j \sqrt{-1}/d)$ is a root of $\lambda_\ell(t) = 0$. Combining these results, we observe that $\lambda_j(t)$ has no multiple roots. Furthermore Libgober and Loeser-Vaqu   showed that

Lemma 1. ([3],[4]) *The polynomial $\Delta(t)$ is written as the product*

$$\tilde{\Delta}(t) = \prod_{k=1}^{d-1} \Delta_k(t)^{\ell_k}, \quad k = 1, \dots, d-1$$

where

$$\Delta_k = (t - \exp(\frac{2k\pi i}{d}))(t - \exp(\frac{-2k\pi i}{d}))$$

and $\ell_k = \dim H^1(P^2, L^{(k)})$.

For the definition of the sheaf $L^{(k)}$, we refer to [2, 1, 4]. We use the method of Esnault-Artal to compute ℓ_k ([2], [1]). Note that for the case of sextics $d = 6$, $\Delta_5(t) = \Delta_1(t) = t^2 - t + 1$, $\Delta_4(t) = \Delta_2(t) = t^2 + t + 1$ and $\Delta_3(t) = (t + 1)^2$.

Let C be a given plane curve of degree d defined by $f(x, y) = 0$ and let $\Sigma(C)$ be the singular locus of C and let $P \in \Sigma(C)$ be a singular point. Consider an embedded resolution of C ,

$\pi : \tilde{U} \rightarrow U$ and let $E_1, \dots, E_s$ be the exceptional divisors. Let us choose (u, v) be a local coordinate system centered at P and let k_i and m_i be the order of zero of the canonical two form $\pi^*(du \wedge dv)$ and π^*f respectively along the divisor E_i . An ideal $\mathcal{J}_{P,k,d}$ of $\mathcal{O}_P$ is generated by the function germ ϕ such that the pull-back $\pi^*\phi$ vanishes of order at least $-k_i + [km_i/d]$ along E_i . Namely

$$\mathcal{J}_{P,k,d} = \{\phi \in \mathcal{O}_P; (\pi^*\phi) \geq \sum_i (-k_i + [km_i/d])E_i\}$$

Let us consider the canonical homomorphisms induced by the restrictions:

$$\sigma_{k,P} : \mathcal{O}_P \rightarrow \mathcal{O}_P / \mathcal{J}_{P,k,d}, \quad \sigma_k : H^0(\mathbf{P}^2, \mathcal{O}(k-3)) \rightarrow \bigoplus_{P \in \Sigma(C)} \mathcal{O}_P / \mathcal{J}_{P,k,d}$$

where the right side of σ_k is the sum over singular points of C . By [1], the integer ℓ_k is equal to the dimension of the cokernel σ_k .

2.1. Non-degenerate singularity. In general, the computation of the kernel and cokernel of the homomorphism $\sigma_{k,P}$ requires an explicit computation of the resolution. However for the class of non-degenerate singularities, these informations can be obtained easily by a toric resolution. See [15, 8] for the definition of non-degenerate singularities. In fact, we do not need any data in detail for a resolution. Let us assume that (u, v) is a fixed local analytic coordinate such that $P = (0, 0)$ and C is defined by a function germ $f(u, v)$ and the Newton boundary $\Gamma(f)$ is non-degenerate with respect to (u, v) . Let $Q_1, \dots, Q_s$ be the primitive covectors which correspond to the edges $\Delta_1, \dots, \Delta_s$ of $\Gamma(f)$ respectively. That is, $\Delta(Q_i, f) = \Delta_i$. Here we use the same terminology as in [8].

Let $\pi : \tilde{U} \rightarrow U$ is the canonical toric modification and let $\hat{E}(Q_i)$ be the exceptional divisor corresponding to Q_i . Recall that the order of zeros of the canonical two form $\pi^*(du \wedge dv)$ along the divisor $\hat{E}(Q_i)$ is simply given by $|Q_i| - 1$ where $|Q_i| := (p + q)$ for $Q_i = {}^t(p, q)$. See [8], p. 178. For a function germ $g(u, v)$, let $m(g, Q_i)$ be the multiplicity of the pull-back π^*g on $\hat{E}(Q_i)$. This number is equal to $d(Q_i, g)$ in [8], which is the minimal value of the linear function Q_i restricted to the Newton boundary of $g(u, v)$. Then the following criterion is essentially due to Merle-Teissier [5] and it is useful for the computation of the ideal $\mathcal{J}_{P,k,d}$.

Lemma 2. *Take a function germ g . Then g is contained in the ideal $\mathcal{J}_{P,k,d}$ if and only if*

$$(\sharp) \quad m(g, Q_i) \geq \left\lceil \frac{k}{d} m(f, Q_i) \right\rceil - |Q_i| + 1, \quad \text{for } i = 1, \dots, s$$

The ideal $\mathcal{J}_{P,k,d}$ is generated by the monomials satisfying the above condition.

Note that the condition $(\sharp)$ can be checked without constructing an explicit toric compactification.

Proof. Let Σ^* be a regular subdivision of $\Gamma^*(f)$ which we use to construct our toric modification $\pi : \tilde{U} \rightarrow U$. We put $Q_0 = {}^t(1, 0)$ and $Q_{s+1} = {}^t(0, 1)$. Then for Q_0, Q_{s+1} the conditions are satisfied. Take the cone $\text{Cone}(Q_i, Q_{i+1})$ and let $T_{i,1}, \dots, T_{i,\nu_i}$ be the covectors which are

inserted for the regular subdivision of the cone $\text{Cone}(Q_i, Q_{i+1})$. Note that the conditions $(\sharp)$ for Q_i, Q_{i+1} are equivalent to

$$m(Q_j, g) > \frac{k}{d}m(Q_j, f) - |Q_j|, \quad j = i, i+1$$

On the other hand, we can write $T_{i,j} = \alpha_j Q_i + \beta_j Q_{i+1}$ for some positive rational numbers α_j, β_j . We also observe that

$$\begin{aligned} m(T_{i,j}, f) &= \alpha_j m(Q_i, f) + \beta_j m(Q_{i+1}, f) \\ m(T_{i,j}, g) &\geq \alpha_j m(Q_i, g) + \beta_j m(Q_{i+1}, g). \end{aligned}$$

The first equality follows from the property: $\Delta(T_{ij}, f) = \Delta_i \cap \Delta_{i+1}$. As $|T_{i,j}| = \alpha_j |Q_i| + \beta_j |Q_{i+1}|$, we obtain

$$\begin{aligned} m(T_{i,j}, g) &\geq \alpha_j m(Q_i, g) + \beta_j m(Q_{i+1}, g) \\ &> \frac{k}{d}(\alpha_j m(Q_i, f) + \beta_j m(Q_{i+1}, f)) - (\alpha_j |Q_i| + \beta_j |Q_{i+1}|) = \frac{k}{d}m(T_{i,j}, f) - |T_{i,j}| \end{aligned}$$

Thus the condition $(\sharp)$ is satisfied for the covectors T_{ij} . The second assertion is obvious. $\square$

3. ALEXANDER POLYNOMIALS OF SEXTICS WITH SIMPLE SINGULARITIES

3.1. Normal forms of simple singularities. For simple singularities, we use the following normal forms.

$$(1) \quad \left\{ \begin{array}{l} A_n : f(u, v) = v^2 + u^{n+1} + (\text{higher terms}) \\ D_k : f(u, v) = v^2 u + u^{k-1} + (\text{higher terms}) \\ E_6 : f(u, v) = v^3 + u^4 + (\text{higher terms}) \\ E_7 : f(u, v) = v^3 + v u^3 + (\text{higher terms}) \\ E_8 : f(u, v) = v^3 + u^5 + (\text{higher terms}) \end{array} \right.$$

Note that $\deg f(0, v) \geq 3$ for D_j and $\deg f(u, 0) \geq 5$ for E_7 and in fact we can make $f(x, y)$ to be convenient by taking $u \mapsto u + v$ or $v \mapsto v + u^2$. In the above notation, we observe that the line $v = 0$ gives the tangent cone of the respective singularities and the local intersection number with the given curve is strictly greater than the respective multiplicity. In the case of D_k singularity, $v = 0$ is one of the tangent cone. Let L_v be the projective line passing through the origin $O = (0, 0)$ in the coordinate (u, v) and tangent to the smooth curve $v = 0$. We call L_v the *principal tangential direction* of the simple singularity. As L_v is written as $v + a u^2 + (\text{higher terms})$, we have $I(L_v, C; O) \geq 4$ for A_j , $j \geq 3$ or E_6, E_7, E_8 or D_k , $k \geq 5$. Here $I(f, g; P)$ is the local intersection number of $f = 0$ and $g = 0$ at P . Let us consider the canonical homomorphisms defined in the previous section:

$$\sigma_{k,P} : \mathcal{O}_P \rightarrow \mathcal{O}_P / \mathcal{J}_{P,k,6}, \quad \sigma_k : H^0(\mathbf{P}^2, \mathcal{O}(k-3)) \rightarrow \bigoplus_P \mathcal{O}_P / \mathcal{J}_{P,k,6}$$

3.2. Description of local data for simple singularities. We assume hereafter (u, v) is a chosen local coordinate system so that the defining equations are written as in (1). The local data for the simple singularities are described by the following.

Proposition 3. *Assume that (C, P) is a simple singularity defined by a normal form as in (1). For $k \leq 3$, $\rho(P, k) = 0$. For $k = 4$ we have*

$$\begin{aligned} \rho(P, 4) &= 0, \text{ for } (C, P) \cong A_1, \dots, A_4 \\ \rho(P, 4) &= 1, \mathcal{J}_{P,4,6} = \langle u, v \rangle \text{ for } (C, P) \cong A_5, \dots, A_{10}, D_4, \dots, D_9, E_6, E_7, E_8 \\ \rho(P, 4) &= 2, \mathcal{J}_{P,4,6} = \langle u^2, v \rangle \text{ for } (C, P) \cong A_{11}, \dots, A_{16}, D_{10}, \dots, D_{15} \\ \rho(P, 4) &= 3, \mathcal{J}_{P,4,6} = \langle u^3, v \rangle \text{ for } (C, P) \cong A_{17}, \dots, A_{22}, D_{16}, \dots, D_{21} \end{aligned}$$

For $k = 5$, we have $\rho(P, 5) = 0$ for A_1 and

$$\begin{aligned} \rho(P, 5) &= 1, \mathcal{J}_{P,5,6} = \langle u, v \rangle \text{ for } (C, P) \cong A_2, A_3, A_4, D_4, D_5 \\ \rho(P, 5) &= 2, \mathcal{J}_{P,5,6} = \langle u^2, v \rangle \text{ for } (C, P) \cong A_5, A_6, A_7, E_6, E_7, E_8, D_6, D_7, D_8 \\ \rho(P, 5) &= 3, \mathcal{J}_{P,5,6} = \langle u^3, v \rangle \text{ for } (C, P) \cong A_8, A_9, A_{10}, D_9, D_{10}, D_{11} \\ \rho(P, 5) &= 4, \mathcal{J}_{P,5,6} = \langle u^4, v \rangle \text{ for } (C, P) \cong A_{11}, A_{12}, A_{13}, D_{12}, D_{13}, D_{14} \\ \rho(P, 5) &= 5, \mathcal{J}_{P,5,6} = \langle u^5, v \rangle \text{ for } (C, P) \cong A_{14}, A_{15}, A_{16}, D_{15}, D_{16}, D_{17} \\ \rho(P, 5) &= 6, \mathcal{J}_{P,5,6} = \langle u^6, v \rangle \text{ for } (C, P) \cong A_{17}, A_{18}, A_{19}, D_{18}, D_{19}, D_{20} \end{aligned}$$

We call a singularity (C, P) is $\rho(k)$ -essential if $\rho(k, P) > 0$. Thus we see that $A_1, \dots, A_4$ are not $\rho(4)$ -essential and A_1 is not $\rho(5)$ -essential.

Corollary 4. I. *Take a germ $g(x, y) \in \mathcal{O}_P$. Then g is contained in the kernel of $\sigma_{k,P}$ if and only if $I(g, v; P) \geq \rho(P, k)$. This implies also*

$$(2) \quad I(g, f; P) \geq 2\rho(P, k)$$

For $k = 4$, the equality in (2) holds only if $(C, P) \cong A_j$, $j \geq 5$.

For $k = 5$, the equality in (2) holds only if either $(C, P) \cong A_j$, $j \geq 2$ or $(C, P) \cong E_6$ singularities. If $(C, P) \cong E_6$ and $I(g, f; P) = 4$, the curve $g = 0$ is smooth at P and it is tangent to the tangent cone of (C, P) .

II. Assume that $g, g' \in \text{Ker } \sigma_{k,P}$. Then $I(g, g'; P) \geq \rho(P, k)$.

Proof. Recall that $f = 0$ is the defining polynomial of C . Let (u, v) be a coordinate system which gives the normal form (1). Write $g = v g_1 + u^r g_2$, $g_1, g_2 \in \mathcal{O}_P$ with $g_2(0) \neq 0$. Then g is contained in the ideal $\mathcal{J}_{P,k,6}$ if and only if $r \geq \rho(P, k)$. Using the normal form in (1), we

have

$$(3) \quad I(u, f; P) = \begin{cases} 2 & (C, P) \cong A_j, j \geq 2 \\ 3 & (C, P) \cong E_6, E_7, E_8 \end{cases}$$

$$(4) \quad I(v, f; P) = \begin{cases} j+1 & (C, P) \cong A_j, j \geq 2 \\ 4, & (C, P) \cong E_6 \\ \geq 5 & (C, P) \cong E_7, E_8 \\ j-1 & (C, P) \cong D_j, j \geq 4 \end{cases}$$

Thus if $r \geq \rho(P, k)$, we have

$$\begin{aligned} I(u^r, f; P) &\geq 2\rho(P, k), \quad I(v, f; P) \geq 2\rho(P, 5) \geq 2\rho(P, 4) \\ I(g, C; P) &\geq \min(I(v, C; P), I(u^r, C; P)) \geq 2\rho(P, k) \end{aligned}$$

The equality $I(v, f; P) = 2\rho(P, 5)$ holds only if $(C, P) \cong E_6$ and $g_1(P) \neq 0$.

Assume $k = 4$. Then by the above inequality, we have $I(g, f; P) \geq 2\rho(P, 4)$. The equality $I(g, f; P) = 2\rho(P, 4)$ holds if and only if $(C, P) \cong A_j$ singularity and $r = \rho(P, 4)$.

Now we assume that $k = 5$. Let g be a conic in $\text{Ker } \sigma_{5,P}$, written as $g = v g_1 + u^r g_2$, $g_1, g_2 \in \mathcal{O}_P$. By the above inequalities, we have $I(v, f; P) \geq 2\rho(P, 5)$ and the equality holds only for E_6 . The equality $I(g, f; P) = I(u^r, f; P) = 2\rho(P, 5)$ takes place if and only if $(C, P) = A_j$ and $r = \rho(P, 5)$. Similarly the equality $I(g, f; P) = I(v, f; P) = 2\rho(P, 5)$ if $(C, P) = E_6$ and $g_1(O) \neq 0$. In the last case, $g = 0$ is smooth at P and its tangent line is the tangent cone of (C, P) .

The assertion II is obvious by the description of the ideal $\mathcal{J}_{P,k,6}$. $\square$

Remark 5. Assume that C is of torus type defined by $f_2^3 + f_3^2 = 0$ and $O \in \{f_2 = f_3 = 0\}$. Assume that $(C, O) \cong A_{3r-1}$. Then the cubic $f_3 = 0$ is smooth at O and $I(f_3, f_2; O) = r$. We may assume that $v = f_3$ and $f_2 \equiv a u^r + (\text{higher terms}) \pmod{(v)}$ for some $a \in \mathbf{C}^*$. Thus $f_2 \in \text{Ker } \sigma_{5,O}$. Assume next that (C, O) is an E_6 -singularity. Then the cubic $f_3 = 0$ has a node at O and f_2 is smooth at O and the tangent cone is equal to the tangent space of $f_2 = 0$ and we may assume that $f_2 = v$ ([11]). Thus again $f_2 \in \text{Ker } \sigma_{5,O}$. Thus $f_2 \in \sigma_{5,O}$ if O is an simple inner singularity.

Lemma 6. *Let ℓ be a projective line and let P_1, P_2, P_3 be three points on ℓ (including the infinitely near point cases $P_1 = P_2 \neq P_3$ or $P_1 = P_2 = P_3$). Let $g = 0$ be a conic on $\mathbf{P}^2$ which passes through them. Then ℓ divides g .*

Proof. The assumption for the degenerated cases $P_1 = P_2 \neq P_3$ or $P_1 = P_2 = P_3$ implies $I(g, \ell; P_1) = 2$ or 3 respectively. Thus in any case, $I(g, \ell) \geq 3$. Thus by Bezout theorem, this implies that $\ell \mid g$. $\square$

Definition 7. *Let $P_1, \dots, P_\nu$, $\nu \leq 3$ be $\rho(5)$ -essential simple singularities on a reduced sextic C such that $\sum_{i=1}^\nu \rho(P_i, 5) \geq 3$. We say that the singular points $\{(C, P_i); i = 1, \dots, \nu\}$ are*

colinear if there is a projective line L such that $P_i \in L$ for $i = 1, \dots, \nu$ and one of the following conditions is satisfied.

(1) $\nu = 3$ or (2) $\nu = 2$, $I(L, C; P_1) \geq 4$ and $\rho(P_1, 5) \geq 2$ and L is the principal tangential line at P_1 or (3) $\nu = 1$ and L is the principal tangential line at P_1 , $I(L, C; P) \geq 6$ and $\rho(P_1, 5) \geq 3$.

Note that in either case, we have $\sum_{i=1}^{\nu} I(L, C; P_i) = 6$ by Bezout theorem.

We recall the definition of *sextics of linear type*. Assume that C is a sextics with $3A_5$ or $A_{11} + A_5$ or A_{17} . We say that C is of *linear type* if there is a line $L \subset \mathbf{P}^2$ such that

$$L \cap C = \begin{cases} \{P_1, P_2, P_3\}, I(C, L; P_i) = 2, & (C, P_i) \cong A_5, i = 1, 2, 3 \\ \{P_1, P_2\}, I(C, L; P_1) = 2, I(C, L; P_2) = 4, & (C, P_1) \cong A_5, (C, P_2) \cong A_{11} \\ \{P_1\}, I(C, L; P_1) = 6, & (C, P_1) \cong A_{17} \end{cases}$$

Let ℓ be the linear form defining L . Then we have shown in [6] that C is of torus type $f_3^2 + f_2^3 = 0$ with $f_2 = \ell^2$. Such a torus curve is called *of linear torus type*. For further detail, see [6]. Now we state our main theorem for sextics with simple singularities.

Theorem 8. *Assume that C is a reduced sextics with only simple singularities.*

(A) *The homomorphism $\sigma_4 : H^0(\mathbf{P}^2, \mathcal{O}(1)) \rightarrow \bigoplus_{P \in \Sigma(C)} \mathcal{O}_P / \mathcal{J}_{P,4,6}$ is described as follows.*

(a) *σ_4 is injective if $\rho(4) \geq 4$. The case $\rho(4) > 4$ does not exist.*

(b) *If $\rho(4) \leq 3$, the homomorphism $\sigma_4 : H^0(\mathbf{P}^2, \mathcal{O}(1)) \rightarrow \bigoplus_{P \in \Sigma(C)} \mathcal{O}_P / \mathcal{J}_{P,4,6}$ is surjective except the case*

(#4) : $\rho(4) = 3$ and C is of linear torus type.

In this case, $\dim \text{Coker } \sigma_4 = 1$.

(B) *The homomorphism $\sigma_5 : H^0(\mathbf{P}^2, \mathcal{O}(2)) \rightarrow \bigoplus_{P \in \Sigma(C)} \mathcal{O}_P / \mathcal{J}_{P,5,6}$ is described as follows.*

(c) *σ_5 is surjective if $\rho(5) \leq 5$.*

(d) *If $\rho(5) \geq 6$, σ_5 is injective except the case that*

(#5) : C is of torus type and $\rho(5) = 6$.

In the last case, $\dim \text{Coker } \sigma_5 = 1$.

The proof of Theorem 8 occupies the rest of this section. Let $P_1, \dots, P_\nu$ be the $\rho(k)$ -essential singularities.

Proof of the assertion (A) in Theorem 8. We first prove the assertion (A). By the classification tables [11, 9, 6] and Bezout theorem, we observe that $\rho(4) \leq 4$ and there exists a unique configuration $\Sigma = [3A_5, D_4]$ which satisfies the equality. Note that the rank is maximal for this configuration.

If there is a line $\ell \in \text{Ker } \sigma_4$, we have $6 \geq I(\ell, C) \geq 2\rho(4)$. This implies $\rho(4) \leq 3$. Thus assume that $\rho(4) \leq 3$. The condition for a linear form ℓ to be in the kernel of σ_4 is given by $\rho(4)$ linear equations. If $\rho(4) < 3$, they are independent by a direct computation. Thus $\dim \text{Ker } \sigma_4 = (3 - \rho(4))$ and the surjectivity follows. Assume that $\rho(4) = 3$ and $\ell \in \text{Ker } \sigma_4$. We may assume that ℓ is not a component of C . For the proof of this assertion, see Appendix 1. By Corollary 4, we have $I(\ell, C) \geq 6$. Thus we must have the equality $I(\ell, C; P) = 2\rho(P)$. Thus the singularities $P \in C$ with $\rho(P, 4) > 0$ must be A_j , $j \geq 5$ and

they are on ℓ so that $I(\ell, C; P) = 2\rho(P, 4)$. As ℓ is unique up to a multiplication of a constant by the property $I(\ell, C) \geq 6$, the assertion (b) reduces to

Lemma 9. *Assume that C is a sextic and ℓ is a line and one of the conditions are satisfied.*

(i) $C \cap \ell = \{P_1, P_2, P_3\}$, $(C, P_i) \cong A_{j(i)}$ with $\rho(P_i, 4) = 1$, $5 \leq j(i) < 11$ and $I(\ell, C; P_i) = 2$ for $i = 1, 2, 3$.

(ii) $C \cap \ell = \{P_1, P_2\}$, $(C, P_i) \cong A_{j(i)}$ with $\rho(P_1, 4) = 2$, $\rho(P_2, 4) = 1$, $11 \leq j(1) < 17$, $5 \leq j(2) < 11$ and $I(\ell, C; P_1) = 4$ and $I(\ell, C; P_2) = 2$.

(iii) $C \cap \ell = \{P_1\}$, $(C, P_1) \cong A_{j(1)}$ with $\rho(P_1, 4) = 3$, $17 \leq j(1) < 23$ and $I(\ell, C; P_1) = 6$. Under the above assumption, we conclude that $j(i) = 6$ or 12 or 18 , namely the singularities on $C \cap \ell$ are $3A_5$ or $A_{11} + A_5$ or A_{17} respectively.

Proof of Lemma 9. Assume that $r_i := \rho(P_i, 4)$ and $(C, P_i) = A_{j(i)}$ with $6r_i - 1 \leq j(i) < 6r_i + 5$. Take normal coordinates (u, v) so that C is defined by $v^2 + u^{j(i)+1} + (\text{higher terms}) = 0$. The assumption $I(\ell, C; P_i) = 2r_i$ implies that $\ell = av + bu^{r_i} + (\text{higher terms})$, $a, b \in \mathbb{C}$ and $b \neq 0$. Let us consider the family of sextics $C_t = f_t^{-1}(0)$, $f_t := f(x, y) + t\ell(x, y)^6$. It is easy to observe that $(C_t, P_i) \cong A_{6r_i-1}$ for a generic $t \in \mathbb{C}$. Thus for such a generic t , C_t is of linear type. Thus by Proposition 7 of [6] or by [14], C_t is of linear torus type written as $f_3(x, y)^2 + c\ell(x, y)^6 = 0$ for some $c \neq 0$. This implies again $C = C_0$ is also of linear torus type, as $f_0 = f_3^2 + (c - \tau)\ell^6$. Now the simple singularities of sextics of linear torus type are $3A_5$, $A_{11} + A_5$ or A_{17} ([6]). $\square$

Proof of the assertion (B) in Theorem 8. Now we consider the assertion (B). Recall that $\dim H^0(\mathbb{P}^2, \mathcal{O}(2)) = 6$. Assume that $\rho(5) < 6$. We will show the surjectivity of σ_5 . Let $P_1, \dots, P_\nu$ be the ρ -essential singularities and let $r_i := \rho(P_i, 5)$. For a conic g , the condition for g to be in the kernel of $\sigma(k, P)$ is given by $\rho(P, k)$ linear conditions. Namely, taking a normal coordinate system (u_i, v_i) of the singularity (C, P_i) , the condition $g \in \text{Ker } \sigma_{5, P_i}$ is given by (considering g as a function of u_i, v_i) $g(0, 0) = \frac{\partial^j g}{(\partial u_i)^j}(0, 0) = 0$, $j = 1, \dots, r_i - 1$ and $i = 1, \dots, \nu$. Thus we have to show that these conditions are independent so that $\dim \text{Image } \sigma_5 = 6 - \dim \text{Ker } \sigma_5 = \rho(5)$. If $\rho(5) \leq 3$, the condition can be checked easily by case-by-case computation. We consider the case $\rho(5) = 4$.

Colinear Case with $\rho(5) = 4$. Assume that $\{(C, P_i); i = 1, \dots, \nu\}$ contains a colinear subset, say $\{(C, P_i); i = 1, \dots, \mu\}$, $\mu \leq \nu$. Let L be a projective line such that $\sum_{i=1}^\mu I(L, C; P_i) = 6$. Then for any conic $g \in \text{Ker } \sigma_5$, we assert $L \mid g$. Otherwise we have a contradiction:

$2 \geq \sum_{i=1}^\mu I(L, g; P_i) \geq 3$. Thus we can write $g = L \times \ell$ with $\ell = ax + by + c$. Now the condition g to be in the kernel of σ_5 is given by one linear condition among a, b, c and therefore the dimension of the kernel σ_5 is 2. For example, consider the easiest case: $\nu = 4$ and $\mu = 3$. Then g is in $\text{Ker } \sigma_5$ if and only if $P_4 \in \ell$. The other possibilities are the cases $\nu = 3, 2, 1$. Consider the case $\nu = \mu = 2$ and $r_1 = r_2 = 2$. We may assume that L is the principal tangent direction of (C, P_1) and P_2 is on L . Note that the principal tangential direction of (C, P_2) is different from L . Then the condition to be asked is $P_2 \in \ell$ as $I(g, C; P_2) \geq 4$. We omit the proof of the other cases as it is similar.

General case with $\rho(5) = 4$. Now we consider the generic case (non-colinear singularities) with $\rho(5) = 4$. The case $\nu = 3$ or 4 or $\nu = 2$ and $r_1 = r_2 = 2$ is easy to be checked, by a direct computation after putting P_i to suitable positions, say for example

$P_1 = (0, 0)$, $P_2 = (1, 0)$, $P_3 = (0, 1)$ and $P_4 = (1, 1)$ for $\nu = 4$ or

$P_1 = (0, 0)$ with the principal tangential line $x = 0$, $P_2 = (1, 1)$, $P_3 = (1, -1)$ for $\nu = 3$ and $r_1 = 2$, etc, using $PGL(3, \mathbf{C})$ -action.

We consider two cases which is slightly less obvious: (1) $\nu = 1$, $r_1 = 4$ or (2) $\nu = 2$, $r_1 = 3$. Assume that (u, v) is a fixed coordinate system which gives the normal form of (C, P_1) . We can choose affine (x, y) coordinates so that $x = x(u, v)$, $y = y(u, v)$ where

$$x(u, v) = u + \sum_{i+j \geq 2} a_{i,j} u^i v^j, \quad y(u, v) = v + \sum_{i+j \geq 2} b_{i,j} u^i v^j$$

Put $P_2 = (\alpha, \beta)$ in the case (2) $\nu = 2$. The genericity condition implies that the principal tangential line $L_v : y = 0$ satisfies $I(L_v, C; P_1) \leq 5$, or equivalently $b_{2,0} \neq 0$ for the case 1 (respectively $P_2 \notin L$, so $\beta \neq 0$ in case 2). Let $g = \sum_{i+j \leq 2} c_{i,j} x^i y^j$ be a conic. We consider the condition $g \in \text{Ker } \sigma_{5, P_1}$ with $P_1 = (0, 0)$. First we need $c_{0,0} = 0$. Let $g_1(u) = g(x(u, 0), y(u, 0))$. By an easy computation, we get

$$\text{Coeff}(g_1, u) = c_{1,0}, \text{Coeff}(g_1, u^2) = c_{0,1}b_{2,0} + c_{2,0}, \text{Coeff}(g_1, u^3) = c_{0,1}b_{3,0} + c_{1,1}b_{2,0} + 2c_{2,0}a_{2,0}$$

Thus in the case 1, we can solve the equation $g_1 \equiv 0$ modulo (u^4) as

$$g(x, y) = c_{0,2} y^2 + \left(\frac{c_{0,1} (-b_{3,0} + 2 b_{2,0} a_{2,0}) x}{b_{2,0}} + c_{0,1} \right) y - c_{0,1} b_{2,0} x^2$$

In the case 2, the condition $g \in \text{Ker } \sigma_5$ is equivalent to $g(\alpha, \beta) = 0$ and $g_1 \equiv 0$ modulo (u^3) which gives

$$g(x, y) = -\frac{(\beta c_{1,1} \alpha + \beta c_{0,1} - c_{0,1} b_{2,0} \alpha^2) y^2}{\beta^2} + (c_{1,1} x + c_{0,1}) y - c_{0,1} b_{2,0} x^2$$

In any case, we see that the kernel is two-dimensional.

Now we consider the case $\rho(5) \geq 5$ and we first show that

Claim 10. *Assume that $\rho(5) \geq 5$, The kernel of σ_5 is at most one-dimensional.*

Proof. Assume first that $\text{Ker } \sigma_5$ contains two mutually coprime conics g, g' . This gives an obvious contradiction

$$4 \geq I(g, g') \geq \sum_{i=1}^{\nu} I(g, g'; P_i) \geq \sum_{i=1}^{\nu} r_i \geq 5$$

Assume next that the kernel contains two linearly independent conics g, g' which has a common linear factor ℓ . Thus we can write $g = \ell \ell_1$ and $g' = \ell \ell'_1$. We may assume that ℓ is not a component of C . See Appendix 2. Put $P := \ell_1 \cap \ell'_1$. Then for any line ξ through P , $\ell \xi$ is again in $\text{Ker } \sigma_5$.

(1) Assume that $P \neq P_i, i = 1, \dots, \nu$. Then taking a generic ξ , the assumption $\ell\xi \in \text{Ker } \sigma_5$ implies that $P_i \in \ell$ for $i = 1, \dots, \nu$. Then we have a contradiction

$$10 \leq 2 \sum_{i=1}^{\nu} r_i \leq \sum_{i=1}^{\nu} I(\ell\xi, C; P_i) = I(\ell, C; P_i) \leq 6$$

(2) Assume that $P = P_1$ for example. The $P_i \in \ell$ for $i = 2, \dots, \nu$. By the assumption, we may assume that ξ is a generic line through P_1 . Thus $I(\xi, C; P_1) \leq 3$. (Recall that the multiplicity of a simple singularity is less than or equal to 3.) By Corollary 4, we have $\sum_{i=1}^{\nu} I(g, C; P_i) \geq 10$. This gives a contradiction that

$$\sum_{i=1}^{\nu} I(\ell, C; P_i) = \sum_{i=1}^{\nu} I(\ell\xi, C; P_i) - I(\xi, C; P_1) \geq 10 - 3 = 7$$

Now the surjectivity of σ_5 for $\rho(5) \leq 5$ is completed.

Assume that $\rho(5) \geq 6$. Assume that there is a conic $g \in \text{Ker } \sigma_5$. We assume that g is not a component of C . See Appendix 3 below for the proof. Then by Corollary 4, we get $12 \geq \sum_{i=1}^{\nu} I(g, C; P_i) \geq 2\rho(5)$ which implies $\rho(5) = 6$. Thus we have $\sum_{i=1}^{\nu} I(g, C; P_i) = 12$ and the conic $g = 0$ intersects C only at ρ -essential singularities. By a result of Tokunaga [13], C is a sextic of torus type. Thus we have proved that σ_5 is injective if $\rho(5) \geq 7$ or $\rho(5) = 6$ and C is not of torus type. This proves the assertion (B) of Theorem 8. $\square$

If C has the rank 19, by Proposition 3 and by the classification table in [16], we see that C is of torus type and C has maximal rank 19. This suggests us:

Conjecture 11. Assume that C is a reduced sextic and assume that $\rho(5) \geq 7$. Then C is a sextic of torus type.

Corollary 12. *Assume that C is a reduced sextic of torus curve with only simple singularities. Then*

1. $\rho(5) = 9$ if and only if C has 9 cusps and in this case $\Delta(t) = (t^2 - t + 1)^3$.
2. Assume that $\rho(5) = 8$.

(a) $\rho(4) \leq 2$ if and only if $\Sigma(C)$ is one of the following.

$$[8A_2], [8A_2, A_1], [6A_2, E_6], [6A_2, A_5], [6A_2, A_5, A_1], [4A_2, 2A_5], [4A_2, A_5, E_6]$$

In this case, C is irreducible and $\Delta(t) = (t^2 - t + 1)^2$.

(b) $\rho(4) \geq 3$ if and only if $\Sigma(C) = [3A_5, 2A_2]$. In this case, C is of linear torus type and C consists of two cuspidal cubic components and $\tilde{\Delta}(t) = (t^2 - t + 1)^2(t^2 + t + 1)$.

3. The case $\rho(5) = 7$ and $\rho(4) = 4$ occurs if and only if $\Sigma(C) = 3A_5 + D_4$. In this case, $\tilde{\Delta}(t) = (t^2 - t + 1)(t^2 + t + 1)$. There are three components in the moduli space:

(a) $C = B_4 + B_1 + B'_1$, where B_4 is a quartic and B_1, B'_1 are two flex tangents of B_4 meeting at a point on B_4 and $\Sigma(C) = [3A_5, D_4]$.

(b) C has three conic components so that $\Sigma(C) = [3A_5, D_4]$.

(c) C is of linear torus with $\Sigma(C) = [3A_5, D_4]$. In this case, C consists of a cubic component and three flex tangent lines.

4. Assume that $\rho(5) \leq 7$ and $\rho(4) = 3$ and C is of linear torus type. Then $\tilde{\Delta}(t) = (t^2 - t + 1)(t^2 + t + 1)$.
5. For other sextics of torus type which is not of linear torus type, we have $\tilde{\Delta}(t) = t^2 - t + 1$. The irreducible sextics of torus type can occur only in 1, 2-(a) or 5.

The proof of Corollary is immediate from Theorem 8 and the classification tables in [11, 9, 6].

3.3. Sextics of non-torus type. We consider irreducible sextics of non-torus type with only simple singularities. We consider the factor $t^2 + t + 1$. This is determined by $\rho(4)$. If $\rho(4) \leq 3$ and C is not of linear torus type, we know by Theorem 8 that the factor $t^2 + t + 1$ does not appear. Now we consider the possibility that $\rho(4) \geq 4$. Recall that the total Milnor number is bounded by 19. Put $\hat{\mu}(4) = \sum_{\rho(P,4)>0} \mu(C, P)$. Proposition 3 implies that $\rho(4) \leq 4$ and the possibility of $\rho(4) = 4$ is the case that $\Sigma(C)$ contains one of the following.

1. $\hat{\mu}(4) = 16$: $4D_4 \subset \Sigma(C)$.
2. $\hat{\mu}(4) = 17$: $3D_4 + X_5 \subset \Sigma(C)$, $X = D, A$.
3. $\hat{\mu}(4) = 18$: $3D_4 + X_6 \subset \Sigma(C)$, $X = D, A, E$ or $2D_4 + X_5 + Y_5 \subset \Sigma(C)$, $X, Y = D, A$ or $D_{10} + 2D_4 \subset \Sigma(C)$.
4. $\hat{\mu}(4) = 19$: $3D_4 + X_7 \subset \Sigma(C)$, $X = D, A, E$ or $2D_4 + X_j + Y_k \subset \Sigma(C)$, $k + j = 11$, $k, j \geq 5$, $X = D, A, E$ or $2D_4 + X_{11} \subset \Sigma(C)$, $X = D, A$, or $D_4 + X_5 + Y_5 + Z_5 \subset \Sigma(C)$, $X, Y, Z = D, A$ or $D_4 + D_{10} + X_5 \subset \Sigma(C)$ with $X = D, A$.

On the other hand, the irreducibility gives the Plücker inequality:

$$\sum_{\rho(P,4)>0} \delta(C, P) \leq 10$$

As $\delta(C, P) = (2\mu(C, P) + r - 1)/2$ where r is the number of irreducible components at P (so $\delta(D_4) = \delta(A_5) = 3$ and so on), we can easily see that none of the above configurations satisfy the Plücker inequality. Thus the factor $t^2 + t + 1$ does not appear in Alexander polynomials of irreducible sextics of non-torus type with simple singularities. By Theorem 8 and Corollary 12 we obtain:

Corollary 13. Assume that C is an irreducible sextics with only simple singularities. Then the Alexander polynomial $\Delta(t)$ takes the form $(t^2 - t + 1)^\alpha$, $\alpha = 0, 1, 2, 3$.

3.4. Appendix. 1. We prove that there does not exist a sextics with simple singularities and $\rho(4) \geq 3$ which has a line component $\ell = 0$ such that $\ell \in \text{Ker } \sigma_4$.

Assume that there exists a sextic $C = L \cup B_5$ with a line component $L = \{\ell = 0\}$ such that $\ell \in \text{Ker } \sigma_4$. Put $P_1, \dots, P_\nu$ be $\rho(4)$ -essential singularities. We assume that (C, P_i) are simple singularities. We will show that $\rho(4) \leq 2$. By the assumption $\ell \in \text{Ker } \sigma_4$ implies that $P_i \in L$ for $i = 1, \dots, \nu$. Note also $\sum_{i=1}^\nu I(L, B_5; P_i) \leq 5$. Thus $\nu \leq 2$.

(1) If $(C, P_i) \cong A_{2k-1} : f(u, v) = v^2 - u^{2k} = 0$, we may assume $\ell = v - u^k$, $I(L, B_5; P_i) = k \leq 5$ and $\rho(4, P) = 1$ for $k = 3, 4, 5$. Actually, $f(u, v)$ may have higher terms and then $\ell = v - u^k + (\text{higher terms})$. We ignored these higher terms also in the following cases but the computation are the same.

(2) If $(C, P_i) \cong E_7 : v^3 + uv^3 = 0$, $\ell = v$, $I(L, B_5; P_i) = 3$ and $\ell \in \text{Ker } \sigma_{5, P_i}$ and $\rho(4, P_i) = 1$.

(3) If $(C, P_i) \cong D_k : v^2u - u^{k-1} = 0$, and $\ell = \{u = 0\}$. Note that $\ell \in \text{Ker } \sigma_{4, P_i}$ if and only if $k \leq 9$. In this case, $I(L, B_5; P_i) = 2$ and $\rho(4, P_i) = 1$. For $k = 2m$, D_{2m} has three components and if $\ell = \{v - u^{m-1} = 0\}$, $I(L, B_5; P_i) = m \leq 5$. Thus $\rho(4, P_i) = 1$ for $m = 2, 3, 4$ and $\rho(4, P_i) = 2$ for $m = 5$.

Thus in any case, we have observed that $\rho(4, P_i) \leq \frac{1}{2}I(\ell, B_5; P_i)$. This implies that

$$\rho(4) = \sum_{i=1}^{\nu} \rho(4, P_i) \leq \sum_{i=1}^{\nu} I(\ell, B_5; P_i) \leq \frac{5}{2}$$

2. Now we show that if ℓ is a component of C and if $\ell\ell_1, \ell\ell'_1$ are linearly independent conics in $\text{Ker } \sigma_5$, then $\rho(5) \leq 4$.

For sextics of torus type, the assertion follows from the list of configurations with a line component in [6]. Assume that C need not be of torus type and ℓ is a component of C and $P_1, \dots, P_\nu$ are on ℓ . Let $C = \ell + B_5$ where B_5 is a reduced curve of degree 5. The possible reducible singularity (C, P_i) is either A_{2k-1} or E_7 or D_j and ℓ must be a smooth component. Let $P = \ell \cap \ell_1$. We can see as before that $P_i \in \ell$ and $\nu \leq 2$ as $\sum_{i=1}^{\nu} I(\ell, B_5; P_i) \leq 5$.

(1) For $A_{2k-1} : v^2 - u^{2k} = 0$, if $\ell = v - u^k$, $I(v - u^k, B_5; O) = k \leq 5$ and $\ell \in \text{Ker } \sigma_{5, P_i}$ and $\rho(5, P_i) \leq [2k/3]$.

(2) For $E_7 : v^3 + vu^3 = 0$, $s \geq 5$, $\ell = v$, $I(L, B_5; O) = 3$ and $\ell \in \text{Ker } \sigma_{5, P_i}$ and $\rho(5, P_i) = 2$.

(3) For $D_k : v^2u - u^{k-1} = 0$,

if (3-1) $\ell = \{u = 0\}$, $I(L, B_5; O) = 2$ and $\rho(5, P_i) = [k/3]$. The assumption $\ell \in \text{Ker } \sigma_{5, P_i}$ implies $k = 4, 5$.

For $k = 2m$ and if (3-2) $\ell = \{v - u^{m-1} = 0\}$, $I(L, B_5; O) = m$ and $\ell \in \text{Ker } \sigma_{5, P_i}$. As $I(L, B_5; P_i) \leq 5$, we need $m \leq 5$.

We observe that in any case,

$$(5) \quad \ell \in \text{Ker } \sigma_{5, P_i} \implies \rho(C, P_i) \leq \left\lceil \frac{2}{3} I(\ell, B_5; P_i) \right\rceil$$

In the above discussion, the higher terms in the defining equations are ignored but the same assertion holds. Put $P = \ell_1 \cap \ell'_1$. We can assume that ℓ_1 is a generic line through P . First, assume that P is not an $\rho(5)$ -essential singularity. This implies that $\ell \in \text{Ker } \sigma_5$. Then we have $\rho(5) \leq \sum_{i=1}^{\nu} \frac{2}{3} I(\ell, B_5; P_i) \leq \frac{10}{3}$ by (5). Next we may assume that $P = P_1$, $(C, P_1) \cong D_k$ and $\ell = u$. The assumption $\ell\ell_1 \in \text{Ker } \sigma_{5, P_1}$ is equivalent to $u^2 \in \text{Ker } \sigma_{5, P_1}$. Thus $\rho(C, P_1) = 2$. The possibilities are $\nu = 2$: $D_i + A_{2j-1}$, ($i = 6, 7, 8$, $j = 2, 3$) or $D_i + E_7$, ($i = 6, 7, 8$) or $\nu = 1$ and D_i , ($i = 6, 7, 8$). Thus in any case we have $\rho(5) \leq 3$.

3. Assume that C is a reduced sextic with only simple singularities. We show that if $g = 0$ a component of a sextics C and $g \in \text{Ker } \sigma_5$, then $\rho(5) \leq 5$.

Assume that $C = C^2 \cup C^4$ where C^2 is defined by $g = 0$ with $g \in \text{Ker } \sigma_5$. Let $P_1, \dots, P_\nu$ be ρ -essential singularities. Then $P_i \in C^2$ for $i = 1, \dots, \nu$. The singularity (C, P_i) is reducible and thus it is one of A_{2j-1} , E_7 or D_j . Note that (5) is true under the substitution $B_5 \rightarrow C^4$. Thus we get

$$\rho(5) = \sum_{i=1}^{\nu} \rho(5, P_i) \leq \frac{2}{3} \sum_{i=1}^{\nu} I(C^2, C^4; P_i) \leq \frac{16}{3}$$

which implies $\rho(5) \leq 5$.

4. NON-SIMPLE SINGULARITIES

In this section, we consider sextics of torus type with some non-simple singularities. By Pho [11], non-simple singularities on sextics of torus type are $B_{3,2k}$ ($k = 2, \dots, 6$), $B_{4,6}, B_{6,6}, C_{3,k}$ ($k = 7, 8, 9, 12, 15$), $C_{6,3k}$ ($k = 2, 3, 4$), Sp_1 , Sp_2 and $D_{4,7}$ where

$$\begin{aligned} B_{m,n} : \quad & f(u, v) = v^m + u^n + (\text{higher terms}) = 0 \quad (\text{Pham-Brieskorn type}) \\ C_{m,n} : \quad & f(u, v) = v^m + u^2 v^2 + u^n + (\text{higher terms}) = 0, \frac{2}{m} + \frac{2}{n} < 1 \\ D_{4,7} : \quad & v^4 + u^3 v^2 + au^5 v + bu^7 = 0, \quad a^2 - 4b \neq 0 \\ Sp_1 : \quad & f(u, v) = f_2(u, v)^3 + f_3(u, v)^2, \quad \begin{cases} f_3(u, v) = v^2 - u^3 + (\text{higher terms}) \\ f_2(u, v) = cuv + (\text{higher terms}), \quad c \neq 0 \end{cases} \\ Sp_2 : \quad & f_3(u, v)^2 - cv^6 = 0, \quad c \neq 0, \quad f_3(u, v) = v^2 - u^3 + (\text{higher terms}) \end{aligned}$$

Note that $B_{m,n}$ and $C_{m,n}$ have non-degenerate Newton boundaries and Sp_1, Sp_2 are degenerated in the sense of Newton boundary ([8]). The following describes the $\rho(k)$ -invariants and the $\mathcal{J}_{P,k,6}$ ideals of the above non-simple singularities. We use the notation $\rho_{3,5}(P) = (\rho(5, P), \rho(4, P), \rho(3, P))$.

Lemma 14. *Assume that C is a sextics and P is a singular point defined by the above equation. Then the ideal $\mathcal{J}_{P,k,6}$ and $\rho(k, P)$ are given as follows.*

1. $B_{3,6}, C_{3,7}$ and $C_{3,8}$:
 $\mathcal{J}_{P,k,6} = \langle u^3, uv, v^2 \rangle, \langle u^2, v \rangle, \langle u, v \rangle$ and $\rho_{3,5}(P) = (4, 2, 1)$.
2. $C_{3,9}$ and $B_{3,8}$: $\mathcal{J}_{P,k,6} = \langle u^4, v^2, uv \rangle, \langle u^2, v \rangle, \langle u, v \rangle$ and $\rho_{3,5}(P) = (5, 2, 1)$.
3. $C_{6,6}$: $\mathcal{J}_{P,k,6} = \langle u^3, v^3, uv \rangle, \langle u^2, v^2, uv \rangle, \langle u, v \rangle$ and $\rho_{3,5}(P) = (5, 3, 1)$.
4. $B_{4,6}, D_{4,7}$: $\mathcal{J}_{P,k,6} = \langle u^3, v^2, vu^2 \rangle, \langle u^2, v^2, uv \rangle, \langle u, v \rangle$ and $\rho_{3,5}(P) = (5, 3, 1)$.
5. $C_{3,12}$: $\mathcal{J}_{P,k,6} = \langle u^5, v^2, uv \rangle, \langle u^2, v \rangle, \langle u, v \rangle$ and $\rho_{3,5}(P) = (6, 2, 1)$.
6. $B_{3,10}$: $\mathcal{J}_{P,k,6} = \langle u^5, v^2, uv \rangle, \langle u^3, v \rangle, \langle u, v \rangle$ and $\rho_{3,5}(P) = (6, 3, 1)$.
7. $C_{6,9}$: $\mathcal{J}_{P,k,6} = \langle u^4, v^3, uv \rangle, \langle u^2, v^2, uv \rangle, \langle u, v \rangle$ and $\rho_{3,5}(P) = (6, 3, 1)$.
8. Sp_1 : $\mathcal{J}_{P,k,6} = \langle u^4, v^2 - u^3, v^3, u^2 v \rangle, \langle u^2, uv, v^2 \rangle, \langle u, v \rangle$ and $\rho_{3,5}(P) = (6, 3, 1)$.
9. $C_{3,15}$: $\mathcal{J}_{P,k,6} = \langle u^6, v^2, uv \rangle, \langle u^3, v \rangle, \langle u, v \rangle$ and $\rho_{3,5}(P) = (7, 3, 1)$.
10. $C_{9,9}$: $\mathcal{J}_{P,k,6} = \langle u^4, v^4, uv \rangle, \langle u^2, v^2, uv \rangle, \langle u, v \rangle$ and $\rho_{3,5}(P) = (7, 3, 1)$.
11. $C_{6,12}$: $\mathcal{J}_{P,k,6} = \langle u^5, v^3, uv \rangle, \langle u^3, v^2, uv \rangle, \langle u, v \rangle$ and $\rho_{3,5}(P) = (7, 4, 1)$.
12. Sp_2 : $\mathcal{J}_{P,k,6} = \langle u^4, v^2 - u^3, v^3, uv^2 \rangle, \langle u^3, uv, v^2 \rangle, \langle u, v \rangle$ and $\rho_{3,5}(P) = (7, 4, 1)$.
13. $B_{3,12}$: $\mathcal{J}_{P,k,6} = \langle u^6, v^2 \rangle, \langle u^4, v \rangle, \langle u^2, v \rangle$ and $\rho_{3,5}(P) = (8, 4, 2)$.
14. $B_{6,6}$: $\mathcal{J}_{P,k,6} = \langle u^4, u^3 v, u^2 v^2, uv^3, v^4 \rangle, \langle u^3, u^2 v, uv^2, v^3 \rangle, \langle u^2, uv, v^2 \rangle, \langle u, v \rangle$ and $\rho_{3,5}(P) = (10, 6, 3)$ and $\rho(2, P) = 1$.

In the above list, $\rho(P, 2) = 0$ except for $B_{6,6}$ and we omitted it.

Proof. Observe that $\mathcal{J}_{P,5,6}$ for Sp_1 and Sp_2 are not monomial ideals. Except for these two singularities, the assertion follows from Lemma 2. We will show the computation for $C_{3,9}$ and

leave the other cases to the reader. Assume that $(C, P) \cong C_{3,9}$. Thus the defining equation is

$$C_{3,9} : f(u, v) = v^3 + u^2v^2 + u^9 = 0$$

We use Lemma 2. We have two faces corresponding to the face function $v^3 + u^2v^2$ and $u^2v^2 + u^9$ whose weight vectors are

$$P_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \quad P_2 = \begin{pmatrix} 2 \\ 7 \end{pmatrix},$$

Let k_i, m_i be the multiplicity of $\pi^*du \wedge dv$ and π^*f along the divisor $\hat{E}(P_i)$. Thus the ideal $\mathcal{J}_{P,k,6}$ is generated by the functions whose pull back have zeros of multiplicity at least $[5m_1/6 - k_1]$ and $[5m_2/6 - k_2]$ along $\hat{E}(P_1)$ and $\hat{E}(P_2)$ respectively. These integers are given by 3 and 7 for $k = 5$ and 2 and 4 for $k = 4$ and 1 and 1 for $k = 3$. On the other hand, $(\pi^*u) = \hat{E}(P_1) + \hat{E}(P_2) + (\text{others})$ and $(\pi^*v) = 2\hat{E}(P_1) + 7\hat{E}(P_2) + (\text{others})$. Thus we can see easily that $\mathcal{J}_{P,k,6}$ is generated by the monomials

$$\mathcal{J}_{P,5,6} = \langle u^4, v^2, uv \rangle, \mathcal{J}_{P,4,6} = \langle u^2, v \rangle, \mathcal{J}_{P,3,6} = \langle u, v \rangle$$

For Sp_1 and Sp_2 , we proceed twice toric modifications to obtain their resolutions.

I. We show the assertion for the case $Sp_1 : f(u, v) = (v^2 - u^3) + c(uv)^3 = 0$. We take the first toric modification $\pi_1 : X_1 \rightarrow \mathbf{C}^2$ with respect to the regular simplicial cone with vertices

$$T_0 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, T_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, T_2 = \begin{pmatrix} 2 \\ 3 \end{pmatrix}, T_3 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, T_4 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

The weight vector T_2 corresponds to the unique edge of the Newton boundary $\Gamma(f; (u, v))$. Consider the toric chart $\text{Cone}(T_2, T_3)$ and denote its toric coordinates by (u_1, v_1) . Then we have

$$u = u_1^2v_1, v = u_1^3v_1, \pi_1^*f(u_1, v_1) = u_1^{12}v_1^6((v_1 - 1)^2 + cv_1^3u_1^3)$$

and the two form $K = du \wedge dv$ is shifted as

$$du \wedge dv = u_1^4v_1^2du_1 \wedge dv_1$$

The strict transform C' of C intersects with $\hat{E}(T_2)$. Take a new coordinate system $(u_1, v'_1) = (u_1, v_1 - 1)$. Now the local behavior of π_1^*f on $\hat{E}(T_2)$ is written as $\pi_1^*(u_1, v'_1) = u_1^{12}(v_1'^2 + cv_1'^3) + (\text{higher terms})$. Thus we proceed the second toric modification $\pi_2 : X_2 \rightarrow X_1$ with respect to the exactly same simplicial cone:

$$S_0 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, S_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, S_2 = \begin{pmatrix} 2 \\ 3 \end{pmatrix}, S_3 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, S_4 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Let (u_2, v_2) be the toric coordinates with respect to $\text{Cone}(S_2, S_3)$. Then $u_1 = u_2^2v_2, v'_1 = u_2^3v_2^2$ and

$$u_1^4du_1 \wedge dv'_1 = u_2^{12}v_2^6du_2 \wedge dv_2$$

Thus putting $\pi = \pi_1 \circ \pi_2 : X_2 \rightarrow \mathbf{C}^2$, we have

$$\begin{aligned} (K) &= \hat{E}(T_1) + 4\hat{E}(T_2) + 2\hat{E}(T_3) + 5\hat{E}(S_1) + 12\hat{E}(S_2) + 6\hat{E}(S_3) \\ (\pi^*f) &= 4\hat{E}(T_1) + 12\hat{E}(T_2) + 6\hat{E}(T_3) + 14\hat{E}(S_1) + 30\hat{E}(S_2) + 15\hat{E}(S_3) \\ (\pi^*u) &= \hat{E}(T_1) + 2\hat{E}(T_2) + \hat{E}(T_3) + 2\hat{E}(S_1) + 4\hat{E}(S_2) + 2\hat{E}(S_3) \\ (\pi^*v) &= \hat{E}(T_1) + 3\hat{E}(T_2) + 2\hat{E}(T_3) + 3\hat{E}(S_1) + 6\hat{E}(S_2) + 3\hat{E}(S_3) \end{aligned}$$

The divisors corresponding to $\mathcal{J}_{P,5,6}$, $\mathcal{J}_{P,4,6}$ and $\mathcal{J}_{P,3,6}$ are given by

$$\begin{aligned} \mathcal{J}_{P,5,6} &= \{\phi; (\phi) \geq 2\hat{E}(T_1) + 6\hat{E}(T_2) + 3\hat{E}(T_3) + 6\hat{E}(S_1) + 13\hat{E}(S_2) + 6\hat{E}(S_3)\} \\ \mathcal{J}_{P,4,6} &= \{\phi; (\phi) \geq \hat{E}(T_1) + 4\hat{E}(T_2) + 2\hat{E}(T_3) + 4\hat{E}(S_1) + 8\hat{E}(S_2) + 4\hat{E}(S_3)\} \\ \mathcal{J}_{P,3,6} &= \{\phi; (\phi) \geq 2\hat{E}(T_2) + 2\hat{E}(S_1) + 3\hat{E}(S_2) + \hat{E}(S_3)\} \end{aligned}$$

Let us consider $\mathcal{J}_{P,5,6}$. It is easy to see that $u^4, u^2v, uv^2, v^3 \in \mathcal{J}_{P,5,6}$. Furthermore we observe that $v^2 - u^3$ is also in the ideal and they generate the ideal. This implies that $\rho(P, 5) = 6$. For $k = 4, 3$, it is easy to see that $\mathcal{J}_{P,4,6} = \langle u^2, uv, v^2 \rangle$ and $\mathcal{J}_{P,3,6} = \langle u, v \rangle$. So $\rho(P, 4) = 3$ and $\rho(P, 3) = 1$. Observe that $\mathcal{J}_{P,5,6}$ is not a monomial ideal.

II. Now we consider Sp_2 defined by $f(u, v) = (v^2 - u^3)^2 + cv^6 = 0$. First we take the same toric modification $\pi_1 : X_1 \rightarrow \mathbf{C}^2$ with respect to the regular simplicial cone with vertices

$$T_0 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, T_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, T_2 = \begin{pmatrix} 2 \\ 3 \end{pmatrix}, T_3 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, T_4 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Then with respect to the toric coordinate (u_1, v_1) of $\text{Cone}(T_2, T_3)$, the pull-back is written as

$$\begin{aligned} \pi^*f(u_1, v_1) &= u_1^{12}v_1^6\{(v_1 - 1)^2 + cu_1^6v_1^2\} \\ &= u_1^{12}\{v_1'^2 + cu_1^6 + (\text{higher terms})\} \end{aligned}$$

Thus we need one more toric modification $\pi_2 : X_2 \rightarrow X_1$ with respect to the covectors

$$R_0 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, R_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, R_2 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, R_3 = \begin{pmatrix} 1 \\ 3 \end{pmatrix}, R_4 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

where the divisor $\hat{E}(R_0) = \hat{E}(T_2)$ and $\hat{E}(R_3)$ corresponds to the face $v_1'^2 + cu_1^6$. By a similar computation, we can show that $\mathcal{J}_{P,5,6} = \langle u^4, v^3, uv^2, u^3 - v^2 \rangle$ and $\mathcal{J}_{P,4,6} = \langle u^3, v^2, uv \rangle$ and $\mathcal{J}_{P,3,6} = \langle u, v \rangle$. This implies $\rho(P, 5) = 7, \rho(P, 4) = 4$ and $\rho(3) = 1$. This completes the proof of Lemma 14.

5. ALEXANDER POLYNOMIAL FOR SEXTICS OF TORUS TYPE

Lemma 15. *Assume that C is a sextic of torus type and (C, P) is a non-simple singularity. Then $I(g, C; P) > 2\rho(k, P)$ for $g \in \mathcal{O}_P \cap \mathcal{J}_{P,k,6}$.*

Proof. The proof follows from the ideal $\mathcal{J}_{P,k,6}$ description given by Lemma 14. We observe that $\rho(5) \geq 7$ for sextics of torus type with at least one non-simple singularity (see [11, 9, 6]). For example, consider the case $B_{3,6} \in C$ and assume that $(C, P) \cong B_{3,6}$. Then $\rho(P, 5) = 4$ and $\rho(P, 4) = 2$. Assume that $g \in \mathcal{O}_P \cap \mathcal{J}_{P,k,6}$. For $k = 5$, we see that $g(u, v) = u^3a(u, v) + v^2b(u, v) + uv c(u, v)$ with some $a, b, c \in \mathcal{O}_P$. As $I(u^3, C; P) = 9, I(v^2, C; P) = 12$ and

$I(uv, C; P) = 9$, we get $I(g, C; P) \geq 9$. For $k = 4$, we can write $g(u, v) = u^2a(u, v) + vb(u, v)$. This implies that $I(g, C; P) \geq 6$. This proves the assertion. For other singularities, similar argument using the linear combination of generators of $\mathcal{J}_{P,5,6}$ and the normal form of the singularities proves the assertion. $\square$

Now the assertion corresponding to Theorem 8 takes the following form.

Theorem 16. *Assume that C is a reduced sextics of torus type with at least one non-simple singularities.*

- (A) *The homomorphism $\sigma_5 : H^0(\mathbf{P}^2, \mathcal{O}(2)) \rightarrow \bigoplus_{P \in \Sigma(C)} \mathcal{O}_P / \mathcal{J}_{P,5,6}$ is injective.*
- (B) (a) *The homomorphism $\sigma_4 : H^0(\mathbf{P}^2, \mathcal{O}(1)) \rightarrow \bigoplus_{P \in \Sigma(C)} \mathcal{O}_P / \mathcal{J}_{P,4,6}$ is injective for $\rho(4) \geq 4$.*
 (b) *If $\rho(4) \leq 3$, the homomorphism $\sigma_4 : H^0(\mathbf{P}^2, \mathcal{O}(1)) \rightarrow \bigoplus_{P \in \Sigma(C)} \mathcal{O}_P / \mathcal{J}_{P,4,6}$ is surjective.*

Proof. Let $P_1, \dots, P_\nu$ be the ρ -essential singularities. We first prove the assertion (A). Assume that g is a conic in $\text{Ker } \sigma_5$. By the classification tables in [11, 9, 6], we see that $\rho(5) \geq 7$. Thus by Corollary 4 and Lemma 15, we get $I(g, C) \geq 2\rho(5) \geq 14$, a contradiction to Bezout theorem if $g = 0$ is not a components of C . Now assume that $g = 0$ is a conic component C^2 of C . Then $C = C^2 + C^4$ where C^4 is a reduced quartic. Assume that P_1 is a non-simple singularity. Then by th assumption $g \in \text{Ker } \sigma_5$, $P_2 \in C^2 \cap C^4$. By Lemma 14, we can see that no smooth component is in $\mathcal{J}_{P_1,5,6}$. Thus $(C^2, P_1) \cong A_1$. In particular, C^2 is a union of two lines. The non-simple singularities which can have two smooth components are $C_{6,6}$, $C_{6,12}$ or $B_{6,6}$. Assume that $(C, P_1) \cong C_{6,6}$. By the classification of reduced sextics with $C_{6,6}$ ([6]), possible configurations are (a) $\Sigma(C) = C_{6,6} + 2A_2 + 2A_1$ and C has two line components and a quartic B_4 or (b) $\Sigma(C) = C_{6,6} + A_5 + 2A_1$ and C has two line components and two conic components. In the case of (a), two A_2 are on the quartic. In case (b), A_5 has to be on the intersection of two conics. In any case, g can not be in $\text{Ker } \sigma_5$. Assume that $(C, P_1) \cong C_{6,12}$. By [6], there are no possibility of sextics with two linear components. Assume that $(C, P_1) \cong B_{6,6}$. This implies that C consists of 6 lines meeting at P_1 . By Lemma 14, no conic can be contained in $\mathcal{J}_{P_1,5,6}$. Thus $g \notin \text{Ker } \sigma_5$.

Now we prove the assertion (B). First we observe that $\rho(P, 4) \geq 2$ for any non-simple singularities which appears on sextics of torus type. Assume that $\rho(4) = 2$ and assume that ℓ, ℓ' be independent linear forms in $\text{Ker } \sigma_4$. Then this gives an contradiction $I(\ell, \ell') \geq 2$. Now we assume that $\rho(4) \geq 3$ and we show that σ_4 is injective. First we assume that ℓ is a line which is not a component of C . Then ℓ can not be in $\text{Ker } \sigma_4$ as otherwise we have a contradiction:

$$6 \geq \sum_{P \in \ell} I(\ell, C; P) > 2\rho(4) \geq 6$$

Now we prove that C does not have a line component ℓ which is in $\text{Ker } \sigma_4$. Assume that P is a non-simple singularity and assume that $\ell = 0$ is a line component such that $\ell \in \text{Ker } \sigma_{P,4}$. Put $C = B_1 + C^5$ where $B_1 = \{\ell = 0\}$ and C^5 is the union of other components. The possibility for (C, P) with $\ell \in \text{Ker } \sigma_{P,4}$ is, by Lemma 14, one of $B_{3,6}, C_{3,k}, B_{3,10}, B_{3,12}$. By [6], $B_{3,10}$ does not appear on reduced sextics and $B_{3,12}$ is only possible for sextics with three conic components. For $B_{3,6}$ or $C_{3,k}$, $\rho(P, 4) = 2$ and we must have other singularity P' with

$\rho(P', 4) > 0$. However by [6], we know that $I(B_1, C^5; P) = 4$ or 5 . Thus B_1 can intersect C^5 at most one point outside of P and thus we have at most $(C, P') \cong A_1$. This is a contradiction to $\rho(P', 4) > 0$. $\square$

Corollary 17. *Assume that C is a reduced sextic of torus type with (at least) a non-simple singularity.*

I. *Assume that C is irreducible. Then*

1. $\Delta(t) = (t^2 - t + 1)^2$ if $\Sigma(C)$ is either $[C_{3,9}, 3A_2]$ or $[B_{3,6}, 4A_2]$.
2. $\Delta(t) = (t^2 - t + 1)$ for other irreducible sextics.

II. *Assume that C is not irreducible. Then we have the following possibilities.*

1. $\tilde{\Delta}(t) = (t^2 - t + 1)^2$ if $\Sigma(C)$ is either $[B_{3,6}, 4A_2, A_1]$ or $[C_{3,9}, 3A_2, A_1]$.
2. $\tilde{\Delta}(t) = (t^2 - t + 1)^2(t^2 + t + 1)$ if $\Sigma(C) = [2B_{3,6}]$.
3. $\tilde{\Delta}(t) = (t^2 - t + 1)^2(t^2 + t + 1)(t + 1)^2$ if $\Sigma(C) = [B_{3,12}]$
4. $\tilde{\Delta}(t) = (t^2 - t + 1)(t^2 + t + 1)$ if $\Sigma(C)$ is one of the following.

$$[C_{6,6}, A_5], [C_{6,6}, A_5, A_1], [C_{6,6}, A_5, 2A_1], [C_{6,12}], [C_{6,12}, A_1], [B_{4,6}, A_5], [D_{4,7}, A_5], [Sp_2]$$

5. $\tilde{\Delta}(t) = (t^2 - t + 1)^4(t^2 + t + 1)^4(t + 1)^4$ if $\Sigma(C) = [B_{6,6}]$.
6. $\tilde{\Delta}(t) = t^2 - t + 1$ for other reduced sextics of torus curve.

The proof of Corollary is immediate from Theorem 16 and the classification tables in [11, 9, 6]. We remark that $\rho(2) \neq 0$ only for the configuration $[B_{6,6}]$.

In [10], it has been observed that a tame sextics C of torus type with $\Sigma(C) = [3A_2, C_{3,9}]$ is exceptional among tame irreducible sextics of torus type in the sense that $\pi_1(\mathbf{P}^2 - C)$ is not isomorphic to $\mathbf{Z}_2 * \mathbf{Z}_3$ and its Alexander polynomial is given by $(t^2 - t + 1)^2$.

REFERENCES

- [1] E. Artal. Sur les couples des Zariski. *J. Algebraic Geometry*, 3:223–247, 1994.
- [2] H. Esnault. Fibre de Milnor d'un cône sur une courbe plane singulière. *Invent. Math.*, 68(3):477–496, 1982.
- [3] A. Libgober. Alexander invariants of plane algebraic curves. In *Singularities, Part 2 (Arcata, Calif., 1981)*, pages 135–143. Amer. Math. Soc., Providence, RI, 1983.
- [4] F. Loeser and M. Vaquié. Le polynôme d'Alexander d'une courbe plane projective. *Topology*, 29(2):163–173, 1990.
- [5] M. Merle and B. Teissier. Conditions d'adjonction, d'après Du Val. In *Séminaire sur les singularités des surfaces*, volume 777 of *Lecture Notes in Math.*, pages 230–245. Springer-Verlag, 1977.
- [6] M. Oka. Geometry of reduced sextics of torus type. *math.AG/0203034*.
- [7] M. Oka. Two transforms of plane curves and their fundamental groups. *J. Math. Sci. Univ. Tokyo*, 3:399–443, 1996.
- [8] M. Oka. *Non-degenerate complete intersection singularity*. Hermann, Paris, 1997.
- [9] M. Oka and D. Pho. Classification of sextics of torus type. *math.AG/0201035*.
- [10] M. Oka and D. Pho. Fundamental group of sextic of torus type. In A. Libgober and M. Tibar, editors, *Trends in Singularities*, pages 151–180. Birkhäuser, Basel, 2002.
- [11] D. T. Pho. Classification of singularities on torus curves of type $(2, 3)$. *Kodai Math. J.*, 24(2):259–284, 2001.
- [12] R. Randell. Milnor fibers and Alexander polynomials of plane curves. In *Singularities, Part 2 (Arcata, Calif., 1981)*, pages 415–419. Amer. Math. Soc., Providence, RI, 1983.

- [13] H.-o. Tokunaga. (2,3) torus sextics and the Albanese images of 6-fold cyclic multiple planes. *Kodai Math. J.*, 22(2):222–242, 1999.
- [14] H.-o. Tokunaga. Galois covers for $\mathfrak{S}_4$ and $\mathfrak{A}_4$ and their applications. *preprint*, 2000.
- [15] A. N. Varchenko. Zeta-function of monodromy and Newton’s diagram. *Invent. Math.*, 37:253–262, 1976.
- [16] J.-G. Yang. Sextic curves with simple singularities. *Tohoku Math. J.*, 48(2):203–227, 1996.

DEPARTMENT OF MATHEMATICS
TOKYO METROPOLITAN UNIVERSITY
1-1 MIMAMI-OHSAWA, HACHIOJI-SHI
TOKYO 192-0397
E-mail: oka@comp.metro-u.ac.jp